

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/308833025>

Doing something about physician burnout

Article *in* The Lancet · September 2016

DOI: 10.1016/S0140-6736(16)31332-0

CITATIONS

0

READS

190

2 authors:



Ronald M Epstein

University of Rochester

218 PUBLICATIONS 12,494 CITATIONS

[SEE PROFILE](#)



Michael R Privitera

University of Rochester

15 PUBLICATIONS 233 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



ATTENDING:MEDICINE, MINDFULNESS AND HUMANITY [View project](#)



Organizational contributions to clinician burnout and workplace violence [View project](#)

All content following this page was uploaded by [Michael R Privitera](#) on 05 October 2016.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.

Doing something about physician burnout



Burnout affects more than half of practising physicians and is on the rise.¹ When burnout was seen as a crisis of wellbeing—affecting physicians' personal lives and work satisfaction—it garnered little public sympathy and could be dismissed as the whining of a privileged class. Now that evidence suggests that burnout negatively affects physicians' effectiveness and availability to patients, as well as patient safety, physicians, health-care organisations, and the public are justifiably worried about quality of patient care and the health of health-care institutions.^{2,3}

According to psychologist Christina Maslach, burnout is when physicians feel emotionally exhausted, depersonalised (ie, cynical and detached), and ineffective; when they feel “an erosion in values, dignity, spirit, and will”.⁴ Burnout is not an expected reaction to hard work—deeply satisfying work can involve tremendous personal sacrifice. It is not the same as depression, yet severe burnout can evolve into it. Initially, burnout might coexist with empathy, satisfaction, and caring, but unrelieved work stress for months diminishes the ability to call forth and sustain those attributes. Although burnout is not new, recent increases are probably due to interactions between individual and organisational factors,⁵ which contribute to a high burden of responsibility, low perceived control, discordance between individual and organisational values, unsupportive work environments, isolation, and loss of meaning.^{6,7} Burnout is not an acute self-limited illness and it has an uncertain prognosis.

In *The Lancet*, Colin West and colleagues⁸ report a comprehensive meta-analysis that raises fundamental questions about what clinicians and health-care organisations should be doing about burnout now. They reviewed 2617 articles, of which 15 randomised trials and 37 cohort studies were of sufficient quality; all but three were done in high-income countries. Interventions reduced overall burnout from 54% to 44%, high emotional exhaustion from 38% to 24%, and high depersonalisation from 38% to 34% among participating physicians. West and colleagues⁸ note that individual (eg, mindfulness, discussion, and stress management) and organisational (eg, work environment) interventions produced similarly large improvements in burnout; while diverse, all of these

programmes share the initial step of enhancement of awareness. For example, mindfulness training can help individuals be aware of burnout in its early phases— noting changes in the body (eg, headaches or muscle tension), emotions (eg, irritability or sarcasm), or thoughts (blaming self or others)—before it becomes unmanageable, name it, and accept that it is present.⁹ Equally important is physicians' awareness of their ability to mitigate burnout: resilience, perspective taking, and cognitive reappraisal.^{10,11} Just as individuals can be mindful of their level of burnout and wellbeing, health-care organisations can monitor these levels as quality indicators and disseminate findings to raise collective awareness and resolve.¹²

Treatment of burnout solely as a disease or failure of individual practitioners is unlikely to be effective. Rather, the individual and system drivers of burnout also need to be addressed. Physicians tend to name external causes, such as productivity pressures and loss of control, yet they should pay equal attention to psychological factors. Just as resilience has a neurocognitive fingerprint,¹¹ so might burnout. Cognitive rigidity, difficulty with ambiguity, setting of boundaries, and forgiving oneself seem to be risk factors that can be addressed, helping physicians manifest adaptive qualities.^{9,11} To promote community and shared vision, meaningful discussions among colleagues and community building can help.


Addressing burnout on an individual level will not be enough in the current health-care environment.¹³ Leaders

Published Online

September 28, 2016
[http://dx.doi.org/10.1016/S0140-6736\(16\)31332-0](http://dx.doi.org/10.1016/S0140-6736(16)31332-0)

See Online/Articles

[http://dx.doi.org/10.1016/S0140-6736\(16\)31279-X](http://dx.doi.org/10.1016/S0140-6736(16)31279-X)



at health-care institutions should take a careful look at what promotes joy, effectiveness, and engagement with clinical practice among their staff and muster the resolve to take positive action.¹³ Physicians, disillusioned by the productivity orientation of administrators and absence of affirmation for the values and relationships that sustain their sense of purpose, need enlightened leaders who recognise that medicine is a human endeavour and not an assembly line. Health planners need to be aware of human limitations—physical, cognitive, and emotional—when layering increasing demands on individual physicians. Health-care organisations should move beyond a culture of endurance, which overvalues stoicism and dismisses complaints as signs of weakness, and help clinicians be better at self-care than they are at present. 21st-century patients demand more than in the past, rightfully so, calling for greater interprofessional coordination. Evidence suggests that the more physicians use electronic health records and computerised physician order entry, the more burnt out they become;¹⁴ intelligent design of health information systems is needed. The explosion in regulatory documentation begs for re-examination and debulking. To address the sheer fatigue of completion of documentation, scribes might be an answer. Realignment of workflows and consideration of the potentially avoidable costs attributable to turnover, retraining, recruitment, malpractice, and loss of patient satisfaction can make such efforts fiscally feasible.

Although optimisation of programmes to address burnout will require further research, we should not wait for perfect understanding before acting; too much is at stake. Institutions should take the lead and address

clinician burnout now that West and colleagues⁸ have described effective models that have been implemented successfully.

*Ronald M Epstein, Michael R Privitera

School of Medicine and Dentistry, University of Rochester, Rochester, NY, USA (RME, MRP); and Family Medicine Research, University of Rochester, Rochester, NY 14610, USA (RME)
ronald_epstein@urmc.rochester.edu

We declare no competing interests.

- 1 Shanafelt TD, Hasan O, Dyrbye LN, et al. Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014. *Mayo Clin Proc* 2015; **90**: 1600–13.
- 2 Lyndon A. Burnout among health professionals and its effects on patient safety. February, 2016. <https://psnet.ahrq.gov/perspectives/perspective/190/burnout-among-health-professionals-and-its-effect-on-patient-safety> (accessed July 24, 2016).
- 3 Shannon D. Physician well-being: a powerful way to improve the patient experience. *Physician Exec* 2013; **39**: 6–12.
- 4 Maslach C, Leiter MP. *The truth about burnout*. San Francisco: Jossey-Bass, 1997.
- 5 Cotton P, Hart P. Occupational wellbeing and performance: a review of organisational health research. *Aust Psychol* 2011; **38**: 118–27.
- 6 Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med* 2014; **12**: 573–76.
- 7 Sikka R, Morath JM, Leape L. The quadruple aim: care, health, cost and meaning in work. *BMJ Qual Saf* 2015; **24**: 608–10.
- 8 West CP, Dyrbye LN, Erwin PJ, Shanafelt TD. Interventions to prevent and reduce physician burnout: a systematic review and meta-analysis. *Lancet* 2016; published online Sept 28. [http://dx.doi.org/10.1016/S0140-6736\(16\)31279-X](http://dx.doi.org/10.1016/S0140-6736(16)31279-X).
- 9 Krasner MS, Epstein RM, Beckman H, et al. Association of an educational program in mindful communication with burnout, empathy, and attitudes among primary care physicians. *JAMA* 2009; **302**: 1284–93.
- 10 Epstein RM, Krasner MS. Physician resilience: what it means, why it matters, and how to promote it. *Acad Med* 2013; **88**: 301–03.
- 11 Southwick SM, Charney DS. *Resilience: the science of mastering life's greatest challenges*. New York: Cambridge University Press, 2012.
- 12 Wallace JE, Lemaire JB, Ghali WA. Physician wellness: a missing quality indicator. *Lancet* 2009; **374**: 1714–21.
- 13 Privitera MR, Rosentstein AH, Plessow F, LoCastro TM. Physician burnout and occupational stress: an inconvenient truth with unintended consequences. *J Hosp Admin* 2014; **4**: 27–35.
- 14 Shanafelt TD, Dyrbye LN, Sinsky C, et al. Relationship between clerical burden and characteristics of the electronic environment with physician burnout and professional satisfaction. *Mayo Clin Proc* 2016; **91**: 836–48.